

1. Rex DK, Cutler CS, Lemmel GT, et al. Colonoscopic miss rates of adenomas determined by back-to-back colonoscopies. [Gastroenterology 1997;112:24-8.](#)
2. Van Rijn JC, Reitsma JB, Dekker E, et al. Polyp Miss Rate Determined by Tandem Colonoscopy: A Systemic Review. [Am J Gastroenterol 2006;101:343-50.](#)
3. Heresbach D, Barrioz T, Ponchon T, et al. Miss rate for colorectal neoplastic polyps: a prospective multicenter study of back-to-back video colonoscopies. [Endoscopy 2008;40:284- 90.](#)
4. Pabby A, Schoen RE, Weissfeld JL, et al. Analysis of colorectal cancer occurrence during surveillance colonoscopy in the dietary Polyp Prevention Trial. [Gastrointestinal Endoscopy 2005;61:385-91.](#)
5. Postic G, Lewin D, Wallace MB, et al. Colonoscopic Miss Rates Determined by Direct Comparison of Colonoscopy with Colon Resection Specimens. [Am J Gastroenterol 2002;97:3182-5.](#)
6. Pickhardt PJ, Nugent PA, Mysliwiec PA, et al. Location of adenomas missed by optical colonoscopy. [Annals of Internal Medicine 2004;141:352-60.](#)
7. Hewett DG, Rex DK. Miss rate of right-sided colon examination during colonoscopy defined by retroflexion: an observational study. [Gastrointest Endosc 2011;74:246-52.](#)